

Network Group – Lung

Mesothelioma							
Regimen	Indication	Drugs\HRG					
Cisplatin\Pemetrexed (version 1.0)	<ul style="list-style-type: none"> First line palliative chemotherapy in PS 0 or 1 patients with mesothelioma. 	Day(s)	Drug	Dose	Route	Comments	
		1	Pemetrexed	500mg/m ²	IV	Infusion in 100mLs Sodium Chloride 0.9% over 10 minutes.	
		1	Cisplatin	75mg/m ²	IV	Infusion in 500mLs Sodium Chloride 0.9% over 2 hours.	
		Type	Days	HRG			
		Procurement	1	Formulary			
Delivery	1	Day Case					
		<ul style="list-style-type: none"> Every 21 days for up to 4 – 6 cycles. Chemotherapy should be discontinued after 2 cycles if there is no symptomatic or radiologic improvement or the patient suffers excessive toxicity 					
Mesothelioma							
Regimen	Indication	Drugs\HRG					
Pemetrexed (version 1.0)	<ul style="list-style-type: none"> Second line palliative chemotherapy in PS 2 or 3 patients with mesothelioma. 	Day(s)	Drug	Dose	Route	Comments	
		1	Pemetrexed	500mg/m ²	IV	Infusion in 100mLS Sodium Chloride 0.9% over 10 minutes.	
		Type	Days	HRG			
		Procurement	1	Formulary			
		Delivery	1	Day Case			
		<ul style="list-style-type: none"> Every 21 days for up to 6 cycles. 					

Mesothelioma						
Regimen	Indication	Drugs\HRG				
Vinorelbine (Oral) (version 1.0)	<ul style="list-style-type: none"> Palliative chemotherapy in mesothelioma. 	Day(s)	Drug	Dose	Route	Comments
		1 + 8	Vinorelbine	60 - 80mg/m ² (max 60mg)		
		Type	Days	HRG		
		Procurement	1 & 8	Formulary		
		Delivery	1 & 8	Day Case		
	<ul style="list-style-type: none"> Every 21 days for up to 4 – 6 cycles. Chemotherapy should not be continued after 2 cycles if there is no evidence of symptomatic or radiologic response to treatment 					
Vinorelbine (IV) (version 1.0)	<ul style="list-style-type: none"> Palliative chemotherapy in mesothelioma. 	Day(s)	Drug	Dose	Route	Comments
		1 + 8	Vinorelbine	25 - 30mg/m ² (max 60mg)	IV	Infusion in 50mLs Sodium Chloride 0.9% over 5-10mins - follow national and local guidance.
		Type	Days	HRG		
		Procurement	1 & 8	Formulary		
		Delivery	1 & 8	Day Case		
	<ul style="list-style-type: none"> Every 21 days for up to 4 – 6 cycles. Chemotherapy should not be continued after 2 cycles if there is no evidence of symptomatic or radiologic response to treatment 					

--

Mesothelioma							
Regimen	Indication	Drugs\HRG					
Cisplatin\ Vinorelbine (IV) (version 1.0)	<ul style="list-style-type: none"> First line palliative chemotherapy in PS 0 or 1 patients with mesothelioma, in those not suitable for a pemetrexed base regimen 	Day(s)	Drug	Dose	Route	Comments	
		1	Cisplatin	50mg/m ²	IV	Infusion in 500mLs Sodium Chloride 0.9% over 1 hour	
		1 + 8	Vinorelbine	25mg/m ²	IV	Infusion in 50mLs Sodium Chloride 0.9% over 5-10mins - follow national and local guidance	
		Type	Days	HRG			
		Procurement	1	Formulary			
		Delivery	1	Day Case			
		Delivery	8	Day Case			
		<ul style="list-style-type: none"> Every 21 days for up to 4 cycles. Chemotherapy should not be continued after 2 cycles if there is no evidence of symptomatic or radiologic response to treatment 					